


Lagrangian multiforms on coadjoint orbits

Anup Anand Singh
University of Leeds

Based on
Lett. Math. Phys. 114, 34 (2024)
arXiv: 2307.07339
Vincent Caudrelier, Marta Dell'Atti, AAS



Students in Theoretical and Mathematical Physics
Edinburgh, March 07, '24



UNIVERSITY OF LEEDS

Outline

- I Lagrangian multiforms for finite-dimensional integrable systems } The what and the why
- II Lie dialgebras and Lax equations } Algebraic background
- III Constructing Lagrangian multiforms on coadjoint orbits } New results
- IV Lagrangian multiform for the rational Gaudin model }
- V Ongoing work and future directions } Connections, generalisations

I Lagrangian multiforms for
finite-dimensional integrable systems

Hamiltonians : the traditional approach to integrability

A $2N$ -dimensional Hamiltonian system is (Liouville) integrable if it possesses N independent conserved quantities H_j in Poisson involution, that is,

$$\{H_i, H_j\} = 0.$$

One of the H_i can be taken as the Hamiltonian of interest H .

This gives us the notion of an integrable hierarchy: each H_k can be used to define a dynamical system each with respect to a "time" variable t_k .

Hamiltonians: the traditional approach to integrability

We have a hierarchy of commuting Hamiltonian flows:

$$\underbrace{\{H_j, H_k\} = 0}_{\text{Poisson involutivity of Hamiltonians}} \Rightarrow \underbrace{\left[\frac{\partial}{\partial t_j}, \frac{\partial}{\partial t_k} \right] = 0}_{\text{commutativity of vector fields}}.$$

This implies path-independence in the multi-time (t_1, \dots, t_N) space.

Think not of a single integrable system,
but of the entire hierarchy it lives in.



Takeaway
message

But how would one capture integrable hierarchies
in the Lagrangian formalism?

Lagrangian multiforms: a variational criterion for integrability

A variational criterion for integrability was introduced in [Lobb-Nijhoff '09] in a discrete setup.

What we need is a collection of Lagrangians \mathcal{L}_k associated with times t_k assembled into a 1-form

$$\mathcal{L}[q] = \sum_{k=1}^N \mathcal{L}_k[q] dt_k.$$

central objects in the Lagrangian multiform theory for finite-dimensional integrable systems

Here q denotes generic configuration coordinates. By $\mathcal{L}[q]$ and $\mathcal{L}_k[q]$, we mean that these quantities depend on q and a finite number of derivatives of q with respects to the times t_1, \dots, t_N .

Lagrangian multiforms: a variational criterion for integrability

We now have an associated generalised action

$$S[q, \Gamma] = \int_{\Gamma} \mathcal{L}[q]$$

this replaces the traditional action
 $S[q] = \int_a^b L[q] dt$

where Γ is a curve in the multi-time \mathbb{R}^N with coordinates t_1, \dots, t_N .

Applying the generalised variational principle to \mathcal{L} gives the multi-time Euler-Lagrange equations

$$\frac{\partial \mathcal{L}_k}{\partial q} - \partial_{t_k} \frac{\partial \mathcal{L}_k}{\partial q_{t_k}} = 0,$$

standard Euler-Lagrange equation for each \mathcal{L}_k

New (corner) Euler-Lagrange equations

$$\frac{\partial \mathcal{L}_k}{\partial q_{t_l}} = 0, \quad l \neq k,$$

Lagrangian coefficient \mathcal{L}_k cannot depend on velocities q_{t_l} for $l \neq k$

$$\frac{\partial \mathcal{L}_k}{\partial q_{t_k}} = \frac{\partial \mathcal{L}_l}{\partial q_{t_l}}, \quad k, l = 1, \dots, N.$$

conjugate momentum to q is the same with respect to all times t_k

Lagrangian multiforms: a variational criterion for integrability

On the solutions of the multi-time Euler-Lagrange equations, we require

$$S[q, \Gamma] = S[q, \Gamma']$$

for all curves Γ, Γ' in the multi-time space.

This implies the closure relation

$$d\mathcal{L}[q] = 0 \iff \partial_{t_k} \mathcal{L}_j - \partial_{t_j} \mathcal{L}_k = 0$$

equivalent to the
Poisson involutivity
of Hamiltonians

on shell.

Lagrangian multiforms: a variational criterion for integrability

These ideas have been extended and illustrated in various other setups:

continuous finite-dimensional systems

[Suris '13] [Petrera-Suris '21]

field theories in 1+1 dimensions

[Suris-Vermeeren '16] [Sleigh-Nijhoff-Caudrelier '19 '20]

[Caudrelier-Stoppato '20 '21] [Petrera-Vermeeren '21]

[Caudrelier-Stoppato-Vicedo '22]

field theories in 2+1 dimensions

[Sleigh-Nijhoff-Caudrelier '21] [Nijhoff '23]

semi-discrete systems

[Sleigh-Vermeeren '22]

Is there an efficient way of describing
all the Lagrangian coefficients in one formula?

II Lie algebras and Lax equations

Lie dialgebras

[Semenov-Tian-Shansky '83]

Let \mathfrak{g} be a Lie algebra with a Lie bracket $[\cdot, \cdot]$, and $R: \mathfrak{g} \rightarrow \mathfrak{g}$ be a linear map. If R is a solution of the modified classical Yang-Baxter equation

$$[R(x), R(y)] - R([R(x), y] + [x, R(y)]) = -[x, y], \quad \forall x, y \in \mathfrak{g},$$

then the skew-symmetric bracket

$$[x, y]_R = \frac{1}{2} ([R(x), y] + [x, R(y)])$$

satisfies the Jacobi identity and defines a second Lie algebra structure on \mathfrak{g} . We will denote the corresponding Lie algebra by \mathfrak{g}_R .

The pair $(\mathfrak{g}, \mathfrak{g}_R)$ is called a Lie dialgebra.

not the same
as a Lie bialgebra

Lie dialgebras

We now have another set of adjoint and coadjoint actions. For $\forall X, Y \in \mathfrak{g}, \forall \xi \in \mathfrak{g}^*$, we can define

$$\text{ad}_X^R \cdot Y = [X, Y]_R \quad \text{and} \quad (\text{ad}_X^{R*} \cdot \xi) Y = -\xi(\text{ad}_X^R \cdot Y) = -\xi([X, Y]_R).$$

We also have the following useful relation

$$R_+ - R_- = \text{Id},$$

adjoint action
of \mathfrak{g}_R on \mathfrak{g}

coadjoint action
of \mathfrak{g}_R on \mathfrak{g}^*

where $R_{\pm} = \frac{1}{2} (R \pm \text{Id})$.

Let $\mathfrak{g}_{\pm} = \text{Im } R_{\pm}$ and $X_{\pm} = R_{\pm}(X)$ for $X \in \mathfrak{g}$. One can show that for any element $X \in \mathfrak{g}$, we have a unique decomposition as

$$X = R_+(X) - R_-(X) = X_+ - X_-.$$

Lie dialgebras

Let us denote by G_R the Lie group associated with the Lie algebra \mathfrak{g}_R . The homomorphisms R_{\pm} give rise to Lie group homomorphisms, which allow us to define the multiplication \cdot^R in G_R as

$$g \cdot^R h = (g_+, g_-) \cdot^R (h_+, h_-) = (g_+ h_+, g_- h_-), \quad \forall g, h \in G_R,$$

where $g_{\pm} h_{\pm}$ denotes the product in G .

We have a new set of adjoint and coadjoint actions, those of G_R on \mathfrak{g}_R and \mathfrak{g}^* , which we can denote in the following useful way:

$$\text{Ad}_g^R \cdot X = \text{Ad}_{g_+} \cdot X_+ - \text{Ad}_{g_-} \cdot X_-, \quad \text{and}$$

$$\text{Ad}_g^{R*} \cdot \xi = R_+^* (\text{Ad}_{g_+} \cdot \xi) - R_-^* (\text{Ad}_{g_-} \cdot \xi), \quad \forall X \in \mathfrak{g}_R, \xi \in \mathfrak{g}^*, g \in G_R.$$

Lie-Poisson bracket and coadjoint orbits

Using the second Lie bracket on \mathfrak{g} , we can define an additional Lie-Poisson bracket on \mathfrak{g}^* , for $f, g \in C^\infty(\mathfrak{g}^*)$ and $\xi \in \mathfrak{g}^*$,

$$\{f, g\}_R(\xi) = (\xi, [\nabla f(\xi), \nabla g(\xi)]_R).$$

natural pairing between \mathfrak{g}^* and \mathfrak{g} : $\xi(x) = (\xi, x)$

the original Lie-Poisson bracket on \mathfrak{g}^* reads $\{f, g\}(\xi) = (\xi, [\nabla f(\xi), \nabla g(\xi)])$

Its symplectic leaves are the coadjoint orbits of G_R in \mathfrak{g}^* .

We need one final ingredient: an Ad-invariant nondegenerate symmetric bilinear form \langle, \rangle on \mathfrak{g} .

allows the identification of \mathfrak{g}^* with \mathfrak{g} and of the coadjoint actions with the adjoint actions

Quick aside: Lax pairs

A Lax pair L, M consists of two matrices - functions on the phase space of the system - such that the equations of motion of the system can be written as

$$\frac{dL}{dt} = [M, L].$$

M is a function
of L

L contains all
information on
the initial data

Spectral invariants of L are integrals of motion.

Involutivity theorem and Lax equations

The Ad^* -invariant functions on \mathfrak{g}^* are in involution with respect to $\{, \}_R$. The equation of motion

$$\frac{d}{dt} L = \{L, H\}_R$$

these functions are simply Casimir functions with respect to $\{, \}$

induced by an Ad^* -invariant function H on \mathfrak{g}^* takes the following equivalent forms, for an arbitrary $L \in \mathfrak{g}^*$,

$$\frac{d}{dt} L = \text{ad}_{\nabla H(L)}^{R*} \cdot L = \frac{1}{2} \text{ad}_{R \nabla H(L)}^* \cdot L = \text{ad}_{R_{\pm} \nabla H(L)}^* \cdot L.$$

using $\{, \}$ would have given trivial equations

Involutivity theorem and Lax equations

The Ad-invariant nondegenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ on \mathfrak{g} allows us to rewrite the last equation in the form of a Lax equation

$$\frac{d}{dt} L = [R_{\pm} \nabla H(L), L].$$

So, the natural arena to define our phase space is a coadjoint orbit of G_R in \mathfrak{g}^* ,

$$\mathcal{O}_{\Lambda} = \{ \text{Ad}_{\varphi}^{R*} \cdot \Lambda; \varphi \in G_R \}, \quad \Lambda \in \mathfrak{g}^*.$$

→
this is where the
Lax matrix L lives



Takeaway
message

Special case: the Adler-Kostant-Symes scheme
[Adler '78], [Symes '78], [Kostant '79]

One gets the well-known Adler-Kostant-Symes scheme by fixing Λ to be in \mathfrak{g}_-^* .

This choice results in only the subgroup G_- in $G_{\mathbb{R}} \cong G_+ \times G_-$ playing a role since

$$L = \text{Ad}_{\mathfrak{g}}^{R*} \cdot \Lambda = -R_-^* (\text{Ad}_{\mathfrak{g}_-}^* \cdot \Lambda).$$

Thus, the coadjoint orbit \mathcal{O}_Λ lies in \mathfrak{g}_-^* .

On to the multi-time story now!

Compatible time flows

For any two Ad^* -invariant functions H_1 and H_2 on \mathfrak{g}^* , we have

$$\{H_1, H_2\}_R = 0.$$

This means that if we have a sufficient number of such independent functions, we can define compatible time flows associated with a family of Ad^* -invariant functions H_k , $k = 1, \dots, N$.

We then obtain an integrable hierarchy with equations in Lax form

$$\partial_{t_k} L = [R_{\pm} \nabla H_k(L), L], \quad k = 1, \dots, N.$$

III Constructing Lagrangian multiforms on coadjoint orbits

The general Lagrangian multiform

[Caudrelier - Dell'Atti - Singh '23]

We introduce the Lagrangian 1-form

$$\mathcal{L}[\varphi] = \sum_{k=1}^N \mathcal{L}_k dt_k = \mathcal{K}[\varphi] - \mathcal{H}[\varphi]$$

with kinetic part

$$\mathcal{K}[\varphi] = \sum_{k=1}^N (L, \partial_{t_k} \varphi \cdot R \varphi^{-1}) dt_k, \quad L = \text{Ad}_{\varphi}^{R*} \cdot \underline{\Lambda}, \quad \underline{\varphi} \in G_R,$$

and potential part

$$\mathcal{H}[\varphi] = \sum_{k=1}^N \underline{H_k(L)} dt_k.$$

fixed non-dynamical element of \mathfrak{g}^*
defining the phase space \mathcal{D}_Λ

field containing the
dynamical degrees of
freedom of the system

Ad^* -invariant
functions $H_k \in C^\infty(\mathfrak{g}^*)$

Euler-Lagrange equations = Lax equations // Result I
[Caudrelier-DeI'Atti-Singh '23]

On considering the variation of the Lagrangian 1-form \mathcal{L} ,
we can derive the Euler-Lagrange equations which take the form

$$\partial_{t_k} L = \frac{1}{2} \operatorname{ad}_{R \nabla H_k(L)}^* \cdot L, \quad k=1, \dots, N.$$

Then, on identifying \mathfrak{g}^* with \mathfrak{g} , and ad^* with ad , we get

$$\partial_{t_k} L = [R_{\pm} \nabla H_k(L), L], \quad k=1, \dots, N,$$

which is exactly the Lax equation associated with the Lax matrix L .

Closure relation // Result II

[Caudrelier-DeI'Atti-Singh '23]

Next, we establish the closure relation for the Lagrangian 1-form \mathcal{L} , that is,

$$d\mathcal{L} = 0, \quad \text{on shell,}$$

or equivalently,

$$\partial_{t_j} \mathcal{L}_k - \partial_{t_k} \mathcal{L}_j = 0, \quad \text{on shell.}$$

This is a consequence of the Ad^* -invariance of H and the fact that R is a solution of the modified CYBE.

Closure relation and Poisson involutivity // Result III

[Caudrelier-Dell'Atti-Singh '23]

Further, for the Lagrangian 1-forms in this class, we can prove

$$\frac{\partial \mathcal{L}_k}{\partial t_\ell} - \frac{\partial \mathcal{L}_\ell}{\partial t_k} = \left\{ H_k, H_\ell \right\}_R = 0, \quad \text{on shell,}$$

demonstrating the connection between the closure relation for Lagrangian 1-forms and the involutivity of Hamiltonians.

This is, in fact, a corollary of a deeper structural result proved in [Caudrelier-Dell'Atti-Singh '23].

first established
in [Sun's '13]

IV Lagrangian multiform for the rational Gaudin model

Gaudin models

Gaudin models are a general class of integrable systems associated with quadratic Lie algebras.

Lie algebras with a nondegenerate invariant bilinear form



First introduced in the quantum finite-dimensional setup to describe quantum spin chains.

[Gaudin '76]

Various generalisations are known — corresponding to both finite- and infinite-dimensional algebras; and with rational, elliptic, skew-symmetric and non-skew-symmetric r -matrices.

A large class of non-ultralocal integrable field theories have been shown to be reinterpretations of classical dihedral affine Gaudin models.

[Vicedo '17]

Rational Gaudin model

The Lax matrix of a rational Gaudin model associated with a finite Lie algebra \mathfrak{g} and a set of points $\zeta_r \in \mathbb{C}$ ($r=1, \dots, N$) and the point at infinity is given by

$$L(\lambda) = \sum_{r=1}^N \frac{X_r}{\lambda - \zeta_r} + X_\infty, \quad X_1, \dots, X_N, X_\infty \in \mathfrak{g},$$

 \mathfrak{g} -valued rational function in variable λ

with the corresponding Lax equations

$$\partial_{t_i^r} X_s = \frac{[X_r, X_s]}{\zeta_r - \zeta_s}, \quad s \neq r,$$


$$\partial_{t_i^r} X_r = -\sum_{s \neq r} \frac{[X_r, X_s]}{\zeta_r - \zeta_s} - [X_r, X_\infty], \quad \partial_{t_i^r} X_\infty = 0.$$

Rational Gaudin model

The quadratic Gaudin Hamiltonians are given as

$$H_r = \sum_{s \neq r} \frac{\text{Tr}(X_r X_s)}{\zeta_r - \zeta_s} + \text{Tr}(X_r X_\infty), \quad r = 1, \dots, N.$$

describes long-range
spin-spin interaction



Algebraic setup

We need to choose a suitable Lie algebra and a linear map from the Lie algebra to itself.

Let us fix

$$Q = \{\zeta_1, \dots, \zeta_N, \infty\} \subset \mathbb{CP}^1,$$

a finite set of points in \mathbb{CP}^1 including the point at infinity, and an index set $S = \{1, \dots, N, \infty\}$.

these become the sites of the model

Denote by

$\tilde{\mathcal{Y}}_Q$ the algebra of \mathfrak{g} -valued rational function in the formal variable λ with poles in Q .

this is where the Lax matrix lives

Define the local parameters $\lambda_r = \lambda - \zeta_r$, $\zeta_r \neq \infty$, and $\lambda_\infty = \frac{1}{\lambda}$.

Algebraic setup

Define the direct sum of Lie algebras

$$\tilde{\mathfrak{g}}_{\mathcal{Q}} = \bigoplus_{r \in \mathcal{S}} \mathfrak{g}_r$$

where

$$\mathfrak{g}_r = \mathfrak{g} \otimes \mathbb{C}((\lambda_r))$$

the Lie algebra
we will work with

is the algebra of formal Laurent series in variable λ_r with coefficients in \mathfrak{g} , and Lie bracket

$$[X\lambda_r^i, Y\lambda_r^j] = [X, Y]\lambda_r^{i+j}, \quad X, Y \in \mathfrak{g}.$$

elements of $\tilde{\mathfrak{g}}_{\mathcal{Q}}$ are tuples
 $(X_1(\lambda_1), \dots, X_N(\lambda_N), X_{\infty}(\lambda_{\infty}))$
with $X_1, \dots, X_N, X_{\infty} \in \mathfrak{g}$

Algebraic setup

We can define a vector space decomposition of $\tilde{\mathfrak{g}}_{\mathcal{Q}}$ into Lie subalgebras

$$\tilde{\mathfrak{g}}_{\mathcal{Q}} = \underbrace{\tilde{\mathfrak{g}}_{\mathcal{Q}^+} \oplus \tilde{\mathfrak{g}}_{\mathcal{Q}^-}}_{\text{decomposition}}$$

we will denote by P_{\pm} the projectors determined by this decomposition

with $\tilde{\mathfrak{g}}_{\mathcal{Q}^{\pm}} = \bigoplus_{r \in \mathcal{S}} \mathfrak{g}_{r^{\pm}}$

where

$$\mathfrak{g}_{r^+} = \mathfrak{g} \otimes \mathbb{C}[[\lambda_r]], \quad r \neq \infty, \quad \text{algebra of formal Taylor series in } \lambda_r$$

$$\mathfrak{g}_{\infty^+} = \mathfrak{g} \otimes \lambda_{\infty} \mathbb{C}[[\lambda_{\infty}]], \quad \text{algebra of formal Taylor series in } \lambda_{\infty} \text{ without the constant term}$$

and

$$\mathfrak{g}_{r^-} = \mathfrak{g} \otimes \lambda_r^{-1} \mathbb{C}[\lambda_r^{-1}], \quad r \neq \infty, \quad \text{algebra of polynomials in } \lambda_r^{-1} \text{ without the constant term}$$

$$\mathfrak{g}_{\infty^-} = \mathfrak{g} \otimes \mathbb{C}[\lambda_{\infty}^{-1}], \quad \text{algebra of polynomials in } \lambda_{\infty}^{-1}$$

Algebraic setup

Further, we have an embedding of Lie algebras

$$L_\lambda : \mathcal{F}_Q(\xi) \hookrightarrow \tilde{\mathfrak{g}}_Q, \quad f \mapsto (L_{\lambda_1} f, \dots, L_{\lambda_N} f, L_{\lambda_0} f),$$

which induces the vector space decomposition

$$\tilde{\mathfrak{g}}_Q = \tilde{\mathfrak{g}}_{Q^+} \oplus L_\lambda \mathcal{F}_Q(\xi).$$

maps $f \in \mathcal{F}_0(\xi)$ to the tuple of its Laurent expansion at points $\xi_1, \dots, \xi_N, \xi_0$

We will denote by Π_\pm the projectors corresponding to this decomposition.
not the same as P_\pm

The r-matrix we need is

$$R = \Pi_+ - \Pi_- .$$

we will use it to equip $\tilde{\mathfrak{g}}_Q$ with a dialgebra structure

Algebraic setup

To identify the dual space to $U_{\lambda} \mathcal{Y}_{\mathbb{Q}}(\mathfrak{g})$, we will use the nondegenerate invariant symmetric bilinear form on \mathfrak{g} ,

$$(X, Y) \mapsto \text{Tr}(XY),$$

to define a nondegenerate invariant symmetric bilinear form on $\tilde{\mathfrak{g}}_{\mathbb{Q}}$:

$$\langle X, Y \rangle = \sum_{r \in S} \text{Res}_{\lambda_r=0} \text{Tr}(X_r(\lambda_r) Y_r(\lambda_r)),$$

which induces the decomposition

$$\tilde{\mathfrak{g}}_{\mathbb{Q}}^* = \tilde{\mathfrak{g}}_{\mathbb{Q}-}^* \oplus \tilde{\mathfrak{g}}_{\mathbb{Q}+}^* \cong \tilde{\mathfrak{g}}_{\mathbb{Q}+}^{\perp} \oplus \tilde{\mathfrak{g}}_{\mathbb{Q}-}^{\perp}.$$

Algebraic setup

Both $\tilde{\mathfrak{g}}_{\mathcal{Q}+}$ and $\mathcal{L}_\lambda \mathcal{Y}_{\mathcal{Q}}(\mathfrak{g})$ are (maximally) isotropic to this bilinear form, which allows us to make the identification

elements of this
are those we
need to work with

$$\tilde{\mathfrak{g}}_{\mathcal{Q}+}^* \cong \mathcal{L}_\lambda \mathcal{Y}_{\mathcal{Q}}(\mathfrak{g}).$$

So, coadjoint orbits of $\tilde{G}_{\mathcal{Q}+}$ in $\tilde{\mathfrak{g}}_{\mathcal{Q}+}^*$ will be the phase space where the Lax matrix of (the model lives and where we will describe its dynamics.

elements of $\tilde{G}_{\mathcal{Q}+}$ are of the form

$$\mathcal{Q}_+ = (\mathcal{Q}_{1+}(\lambda_1), \dots, \mathcal{Q}_{N+}(\lambda_N), \mathcal{Q}_{\infty+}(\lambda_\infty))$$

with $\mathcal{Q}_{r+}(\lambda_r) = \sum_{n=0}^{\infty} \phi_r^{(n)} \lambda_r^n$

and $\mathcal{Q}_{\infty+}(\lambda_r) = 1 + \sum_{n=1}^{\infty} \phi_\infty^{(n)} \lambda_\infty^n$

Algebraic setup

The coadjoint orbit of an element $c_\lambda f \in \mathfrak{g}_{\mathcal{Q}_+}^*$ is given by

$$\begin{aligned}c_\lambda F &= \text{Ad}_{\mathcal{Q}}^{R^*} \cdot c_\lambda f \\ &= R_+^* (\text{Ad}_{\mathcal{Q}_+}^* \cdot c_\lambda f) \\ &= R_+^* (\mathcal{Q}_+ \cdot c_\lambda f \cdot \mathcal{Q}_+^{-1}) \\ &= \Pi_- (\mathcal{Q}_+ \cdot c_\lambda f \cdot \mathcal{Q}_+^{-1})\end{aligned}$$

since we are looking at an element from a subspace of the dual only one corresponding subgroup plays a role in the coadjoint orbit

where we have made the identification $R_+^* = \Pi_-$.

So, we are now ready with our setup!

Lax matrix

Choose

$$\Lambda(\lambda) = \sum_{r=1}^N \frac{\Lambda_r}{\lambda - \rho_r} + \Omega, \quad \Lambda_r, \Omega \in \mathfrak{g}$$

and consider its embedding into $\tilde{\mathfrak{g}}_{\mathcal{Q}}$

$$\iota_{\lambda} \Lambda(\lambda) = \iota_{\lambda} \left(\sum_{r=1}^N \frac{\Lambda_r}{\lambda - \rho_r} + \Omega \right) \in \iota_{\lambda} \mathcal{F}_{\mathcal{Q}}(\mathfrak{g}) \simeq \tilde{\mathfrak{g}}_{\mathcal{Q}+}^*$$

The orbit of $\iota_{\lambda} \Lambda$ under the coadjoint action of $\tilde{\mathcal{G}}_{\mathcal{Q}+}$ will be

$$\begin{aligned} \iota_{\lambda} L &= \pi_{-} \left(\mathcal{Q}_{+} \cdot \iota_{\lambda} \Lambda \cdot \mathcal{Q}_{+}^{-1} \right) \\ &= \iota_{\lambda} \left(\sum_{r=1}^N \frac{A_r}{\lambda - \rho_r} + \Omega \right). \end{aligned}$$

$$A_r = \Phi_r^{(0)} \Lambda_r \Phi_r^{(0)-1}$$

contains the dynamical degrees of freedom

fixed non-dynamical element

Lagrangian multiform for the rational Gaudin model

[Caudrelier - Dell'Atti - Singh '23]

We can now write down the Gaudin multiform on the orbit of $\Lambda(\lambda)$, with the elements $L_\lambda L$,

$$\mathcal{L} = \sum_{k=1}^N \sum_{r \in S} \mathcal{L}_{k,r} dt_k^r,$$

with

$$\mathcal{L}_{k,r} = \sum_{s \in S} \operatorname{Res}_{\lambda_s \neq 0} \operatorname{Tr} (L_{\lambda_s} L \partial_{t_k^r} \mathcal{Q}_{s+}(\lambda_s) \mathcal{Q}_{s+}(\lambda_s)^{-1}) - H_{k,r}(L_\lambda L).$$

restriction of $H_{k,r}$ to $L_\lambda L$

Upon simplification, the Lagrangian coefficients take the form

$$\mathcal{L}_{k,r} = \sum_{s=1}^N \operatorname{Tr} (\Lambda_s \phi_s^{-1} \partial_{t_k^r} \phi_s) - H_{k,r}(L_\lambda L).$$

$\hookrightarrow \phi_s^{(0)} = \phi_s$ for notational simplicity

Lagrangian multiform for the rational Gaudin model

[Caudrelier - Dell'Atti - Singh '23]

The potential part $H_{k,r}(L)$ is the restriction to L of invariant functions on $\tilde{\mathfrak{g}}_{\mathbb{Q}}$ that can be given by

$$H_{k,r} : X \in \tilde{\mathfrak{g}}_{\mathbb{Q}} \longmapsto \operatorname{Res}_{\lambda_r=0} \frac{\operatorname{Tr}(X_r(\lambda_r)^{k+1})}{k+1}, \quad k \geq 1.$$

For $k=1, 2$, we have

$$H_{1,r}(L) = \sum_{s \neq r} \frac{\operatorname{Tr}(A_r A_s)}{\varrho_r - \varrho_s} + \operatorname{Tr}(A_r \Omega)$$

and

$$H_{2,r}(L) = \operatorname{Tr} \left(A_r \left(\sum_{s \neq r} \frac{A_s}{\varrho_r - \varrho_s} + \Omega \right)^2 \right) - \operatorname{Tr} \left(A_r^2 \left(\sum_{s \neq r} \frac{A_s}{(\varrho_r - \varrho_s)^2} \right) \right).$$

Euler-Lagrange equations

Varying $\mathcal{L}_{1,r}$ and $\mathcal{L}_{2,r}$ with respect to ϕ_s , $s=1, \dots, N$, gives the Euler-Lagrange equations for the first and the second time flows respectively:

$$\partial_{t_1^r} A_s = \frac{[A_r, A_s]}{\mathcal{I}_r - \mathcal{I}_s}, \quad s \neq r,$$

$$\partial_{t_1^r} A_r = -\sum_{s \neq r} \frac{[A_r, A_s]}{\mathcal{I}_r - \mathcal{I}_s} - [A_r, \Omega],$$

$$\partial_{t_2^r} A_s = \frac{-[A_r^2, A_s]}{(\mathcal{I}_r - \mathcal{I}_s)^2} + \sum_{s' \neq r} \frac{[A_r A_{s'} + A_{s'} A_r, A_s]}{(\mathcal{I}_r - \mathcal{I}_s)(\mathcal{I}_r - \mathcal{I}_{s'})} + \frac{[A_r \Omega + \Omega A_r, A_s]}{\mathcal{I}_r - \mathcal{I}_s}, \quad s \neq r,$$

$$\partial_{t_2^r} A_r = \sum_{s \neq r} \frac{[A_r^2, A_s]}{(\mathcal{I}_r - \mathcal{I}_s)^2} - \sum_{s \neq r} \sum_{s' \neq r} \frac{[A_r, A_s A_{s'}]}{(\mathcal{I}_r - \mathcal{I}_s)(\mathcal{I}_r - \mathcal{I}_{s'})} - \sum_{s \neq r} \frac{[A_r, A_s \Omega + \Omega A_s]}{\mathcal{I}_r - \mathcal{I}_s} - [A_r, \Omega^2].$$

v Ongoing work and future directions

Ongoing work and future directions

I Lagrangian multiform for cyclotomic Gaudin models
[work in progress with V. Caudrelier and B. Vicedo]

Decomposition of a suitable Lie algebra into subalgebras that are not isotropic with respect to the chosen bilinear form - gives a non-skew-symmetric r -matrix

→ using the full power of the Lie dialgebra framework

Realisation of cyclotomic Gaudin models as some interesting finite-dimensional integrable models

Ongoing work and future directions

II Lagrangian multi-form for affine Gaudin models

reinterpretation as
non-ultralocal integrable
field theories

path integral quantisation
of integrable field theories

?

III Connections to geometric actions

IV Connections of the Lie dialgebra construction with the gauge-theoretic approach to integrability, in particular, mixed BF theory-based construction of Gaudin models

A not-at-all exhaustive list of references

- I S. Lobb, F.W. Nijhoff, Lagrangian multiforms and multidimensional consistency, 2009
- II M.A. Semenov-Tian-Shansky, Integrable systems: the r-matrix approach, 2008
- III V. Caudrelier, M. Stoppato, B. Vicedo, Classical Yang-Baxter equation, Lagrangian multiforms and ultralocal integrable hierarchies, 2022
- IV S. Lacroix, Integrable models with twist function and affine Gaudin models, PhD thesis, 2018
- V V. Caudrelier, M. Dell'Atti, A.A. Singh, Lagrangian multiforms on coadjoint orbits for finite-dimensional integrable systems, 2023

Thank you!

